

MYCOTAXON

<http://dx.doi.org/10.5248/130.601>

Volume 130, pp. 601–605

July–September 2015

**Conidial fungi from the semiarid Caatinga biome of Brazil:
a new species of *Selenosporella* from submerged leaves**PATRÍCIA OLIVEIRA FIUZA¹, LUÍS FERNANDO PASCHOLATI GUSMÃO^{1*},
& RAFAEL F. CASTAÑEDA-RUIZ²¹*Departamento de Ciências Biológicas, Laboratório de Micologia, Universidade Estadual de Feira de Santana, Avenida Transnordestina s/n, Novo Horizonte, 44036-900, Feira de Santana, Brazil*²*Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT), Académico Titular de la Academia de Ciencias de Cuba, Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200** CORRESPONDENCE TO: lgusmao@uefs.br

ABSTRACT — *Selenosporella minima* sp. nov., collected on submerged leaves of *Calophyllum brasiliense* in the Brazilian semiarid region, is described and illustrated. It is distinguished by verticillate branched conidiophores and navicular to acrose conidia.

KEY WORDS — asexual fungi, taxonomy, freshwater

Introduction

Selenosporella G. Arnaud ex MacGarvie comprises twelve species. Descriptions, illustrations, and a key to *Selenosporella* species has been provided by Castañeda-Ruiz et al. (2009), who divided the genus into two groups: i) with conidiogenous cells discrete and arranged in verticils and ii) with conidiogenous cells integrated and not arranged in verticils.

Most *Selenosporella* species have been described from litter or wood (Seifert et al. 2011), except for *Selenosporella aristata* Kuthub. & Nawawi, which was described from submerged leaves (Kuthubutheen & Nawawi 1988). An interesting *Selenosporella* specimen recently found on decaying submerged leaves of *Calophyllum brasiliense* is described here as new.

Materials & methods

During several expeditions between November 2013 and May 2014 through Serra da Tromba in the semiarid region of northeastern Brazil, samples of submerged leaves

of *Calophyllum brasiliense* were stored in plastic bags, taken to the laboratory, and treated according to Castañeda-Ruiz (2005). Mounts were prepared in PVL (polyvinyl alcohol, lactic acid, and phenol) and measurements made at a magnification of $\times 1000$. Microphotographs were obtained with an Olympus microscope (model BX51) with bright field and Nomarski interference optics. The holotype is deposited in the Herbarium of Universidade Estadual de Feira de Santana, Bahia, Brazil (HUEFS).

Taxonomy

Selenosporella minima Fiuza, Gusmão & R.F. Castañeda, **sp. nov.**

FIGS 1, 2

IF 550802

Differs from *Selenosporella cymbiformis* by single, verticillate branched conidiophores.

TYPE: Brazil, Bahia, Piatã, Serra da Tromba, 13°05'S 41°50'W, on submerged leaves of *Calophyllum brasiliense* Cambess. (*Calophyllaceae*), 1-V-2014, coll. P.O. Fiuza (Holotype: HUEFS 210422).

ETYMOLOGY: Latin, *minima*, meaning small, refers to conidial size.


COLONIES on the natural substrate effuse, hairy, brown. Mycelium mostly immersed, composed of septate, branched, brown hyphae. CONIDIOPHORES distinct, single, erect, straight, verticillate branched, 2–5-septate, smooth, dark brown, $60\text{--}100 \times 4\text{--}5 \mu\text{m}$. CONIDIOGENOUS CELLS holoblastic, denticulate, sympodially elongated, discrete, brown to pale brown. Conidial secession schizolytic. CONIDIA solitary, navicular to acerose, unicellular, smooth, hyaline, $3.5\text{--}5 \times 1 \mu\text{m}$.

NOTE: *Selenosporella cymbiformis* B. Sutton is similar to *S. minima* in conidial shape and size, but *S. cymbiformis* has single or loosely synnematus, unbranched or irregularly branched, $35\text{--}55 \times 3\text{--}4 \mu\text{m}$, dark to medium brown conidiophores (Sutton 1973).

Acknowledgments

The authors express their sincere gratitude to Dr. De-Wei Li, Prof. Dr. Meng Zhang, and Dr. Shambhu Kumar, for their critical review of the manuscript. The authors are grateful to the “Programa de Pesquisa em Biodiversidade” – (PPBio Semi-arid/MCTI/CNPq) for financial support. POF thanks the ‘Coordenação de Aperfeiçoamento de Pessoal de Nível Superior’ (CAPES) and the ‘Programa de Pós-graduação em Botânica PPGBot/UEFS’. The authors thank the support provided by ‘Programa Ciência sem Fronteiras’. RFCR is grateful to and Cuban Ministry of Agriculture and ‘Programa de Salud Animal y Vegetal’, project P131LH003033 for facilities. We acknowledge the facilities provided by Dr. P.M. Kirk and Drs. V. Robert and A. Decock through the

FIG. 1. *Selenosporella minima* (ex holotype, HUEFS 210422): A, B. Conidiogenous cells and conidia; C. Conidiophore, conidiogenous cells and conidia; D, E. Conidiophores. Scale bars: A = 10 μm ; B, C = 5 μm ; D, E = 20 μm .






FIG 2. *Selenosporella minima* (ex holotype, HUEFS 210422):
A. Conidia; B. Conidiogenous cells and conidia; C. Conidiophores.
Scale bars: A, B = 5 μ m; C = 20 μ m.

IndexFungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

Literature cited

- Castañeda Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183, in: Anais do V Congresso Latino Americano de Micología. Brasília.
- Castañeda Ruiz RF, Guerrero B, Adamo GM, Morillo O, Minter DW, Stadler M, Gené J, Guarro J. 2009. A new species of *Selenosporella* and two microfungi recorded from a cloud forest in Mérida, Venezuela. *Mycotaxon* 109: 63–74. <http://dx.doi.org/10.5248/109.63>
- Kuthubutheen AJ, Nawawi A. 1988. A new species of *Selenosporella* (hyphomycetes) from Malaysia. *Trans. Br. Mycol. Soc.* 91: 331–334. [http://dx.doi.org/10.1016/S0007-1536\(88\)80222-5](http://dx.doi.org/10.1016/S0007-1536(88)80222-5)
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.
- Sutton BC. 1973. Hyphomycetes from Manitoba and Saskatchewan, Canada. *Mycol. Pap.* 132. 143 p.